



## Face and Neck Lift Pre-Operative Instructions

### Pre-operative Information for General Anaesthetic procedures

This pre-operative guide aims to help you prepare for your surgery. Please let us know if you have any new medical problems or medications prior to surgery.

#### Smokers

You should cease smoking for a minimum of four weeks before the surgery, although discontinuing earlier is preferable. Nicotine diminishes blood oxygen levels and can markedly elevate the likelihood of post-operative healing complications. Since it's the nicotine and not the smoke itself that poses issues, this caution **extends to nicotine replacement therapies** such as chewing gum, e-cigarettes, and patches.

#### Sickle cell trait or sickle cell disease

If you have either condition, it is essential that we are made aware of this prior to your surgery to ensure that you are optimised and have blood transfusions and anticoagulation if necessary. This will be fully discussed at your initial consultation.

#### Compression garments

You will be advised to purchase a chin strap prior to surgery. If required, my nurse can size you in clinic and advise you as to which garments to purchase. You can either buy these from the Hospital before surgery, or alternatively, we can provide you with the supplier details should you wish to obtain your own.

I would recommend that you obtain 2 garments, as you will want one to wash and one to wear. They need to be worn day and night for 2 weeks after surgery, although you can remove them for washing.

#### Medications

Certain medications should be stopped prior to surgery as they can cause blood clots or excessive bleeding. These include:

- **Aspirin / Clopidogrel:** To be stopped 2 weeks prior to surgery as they can cause excessive bleeding and haematoma formation.
- **Warfarin:** This should be stopped 4 to 5 days before surgery, depending on your INR and the indications for the Warfarin. We will start a blood thinning injection to protect you whilst the Warfarin has been discontinued.



## Face and Neck Lift Pre-Operative Instructions

- **Tamoxifen:** To be stopped this 4 weeks before surgery as Tamoxifen can cause blood clots.
- **Oral Contraceptives/HRT:** Hormonal medications should be stopped 4 weeks before surgery as they can increase the risk of blood clots.

### Pre-operative assessment

Prior to your surgery, you will be given a pre-operative assessment with the anaesthetist. This can be either over the phone or in the clinic, if the anaesthetist wishes to see you face to face prior to surgery. Depending on your age and general health, routine blood tests, MRSA swabs and a heart tracing (ECG) may be taken.

You will also have an opportunity to ask any questions you may have or raise any concerns.

If you are unsure whether you wish to proceed with a procedure, then it is strongly advised not to proceed with surgery.

### Admission to hospital

This will normally be on the day of your operation. If you are having a general anaesthetic, you will be asked not to eat anything 6 hours before surgery. Black tea/ coffee or non-fizzy drinks (nothing containing milk) can be taken up to 4 hours before surgery. Chewing gum should also be avoided for 6 hours before your operation.

Sometimes delays do occur, however, you will be cared for whilst waiting under the direction of the anaesthetist and the nursing team.

I will come and see you with the anaesthetist before you go to the operating room. I will take pre-operative photographs and go over the operation again to remind you of any risks or complications.

If you have any questions or concerns, there will always be sufficient time made to answer them.



Anne Dancey  
PLASTIC SURGERY

## Face and Neck Lift Pre-Operative Instructions

At any time if you are unsure about anything concerning surgery, then please speak to me.



I hope you find this information useful. If you have any questions or require a little more information then please do not hesitate to contact me.

*Anne Dancey*

**Dr Anne Dancey**

Plastic and Reconstructive Surgeon FRCS(Plast), MBChB(Hons), MmedSci(Hons)  
and MCh(PASP)